

## Obsah

|  |     |
|--|-----|
| Investigation of the carbon material based electrodes for supercapacitors  | 9   |
| The effect of ochratoxin A on DNA adduct formation by the carcinogen aristolochic acid in rats   | 12  |
| Role of rat cytochromes p450 in oxidation of 17 $\alpha$ -ethinylestradiol   | 15  |
| Structure and spectroscopic characterization of copper(II) complexes with fluoroquinolones   | 18  |
| NAD(P)H:quinone oxidoreductase 1 is Induced by 3-nitrobenzanthrone, aristolochic acid, ellipticine, Sudan I and dicoumarol in rats - a comparative study | 20  |
| Study of structure and chemical composition of soil humic substances isolated from Humic Podzol  | 23  |
| Effect of bitumen on sorption properties of lignite  | 28  |
| Piezoelectric biosensors for detection of microorganisms   | 32  |
| High performance sulfur nanocomposites as cathode materials based on conversion reaction   | 26  |
| Physicochemical methods as a suitable instrument for short DNA fragments study   | 40  |
| Studium fotochemických chránících skupin pomocí časově rozlišené spektroskopie. Je nano sekunda málo nebo moc pro roztržení vazby?                       | 43  |
| Hyaluronana micro- and nanoparticles   | 44  |
| Mircorheological characterization of hyaluronic acid gels  | 47  |
| Development of the cell systém for evaluation of ANTI-PAIIL immunoglobulin efficacy  | 51  |
| Cytochrome p450 1A1-catalyzed oxidation of carcinogenic benzo[a]pyrene is modulated by NADH and cytochrome b <sub>5</sub>                                | 54  |
| The azobenzene actinometer   | 57  |
| Deeper Research of Structural Changes of Humic Acids Using Light Scattering Techniques   | 59  |
| Are flavonoids fenoxide anions better hydrogen atom donors than parent molecules?  | 64  |
| On the homolytic and heterolytic O–H bond cleavage in vitamin b6   | 67  |
| Observation of prompt ISC from higher excited states of the anion of p-hydroxyacetophenone   | 70  |
| Acidity of Frozen Solutions and Its Connection to Degradation of Enzymes upon Freezing   | 71  |
| Study of nanoparticles formed by negatively charged hyaluroan and cationic surfactant  | 73  |
| Application of CdTe-Quantum dots nanoparticles for luminiscence determinativ of metal ions   | 78  |
| Nanosecond laser flash photolysis Study of rose Bengal   | 81  |
| EPR-UV/Vis/NIR SpectroElectrochemical studies of substituted diarylaminothiophenes   | 82  |
| EPR-UV/Vis/NIR SpectroElectrochemistry of thienyl derivatives of pyrene  | 85  |
| How to measure quantitative EPR spectra reproducibly   | 88  |
| Utilization of fluorescence spectroscopy for studying of the interactions in dried systems of biopolymer and hydrophobic species                         | 91  |
| Determination of organic acids in wine using by biosensors   | 95  |
| New Kinetic Models in Biopolymer Degradation: Long Term Study of Hyaluronan Samples  | 99  |
| Oxidation of and DNA adduct formation by benzo[a]pyrene by rat cytochrome p450 1A1 are stimulated by cytochrome b <sub>5</sub> and NADH                  | 103 |

|   |     |
|---|-----|
| Aryl hydrocarbon receptor ligands benzo[a]pyrene, ellipticine and Sudan I induce expression of cytochrome p450 1A1, NAD(P)H:quinone oxidoreductase 1 and cytochrome b <sub>5</sub>                  | 107 |
| Determination of methylxantines on pencil graphite electrodes by cyclic voltammetry   | 111 |
| Trends in development of functional nanostructured films  | 114 |
| Biodegradable materials for orthopedic applications   | 118 |
| An Electrochemical study of d(GCGAAGC) DNA heptamer and its sequential analogues  | 121 |
| Aromatase (Cytochrome p450 19) is an efficient enzyme activating anticancer drug ellipticine  | 124 |
| DFT and NMR studies of PPD antioxidants   | 128 |
| Computational Insight to the Thermodynamics of Double H atom abstraction in model phenolics   | 131 |
| On radical anion generation upon hydrogen atom transfer in dihydroxybenzenes  | 134 |
| Methylation of Humic acids – the Impact on the Reactivity Studied by Diffusion Techniques   | 137 |
| Electrochemical sensor for carbonate determination  | 142 |
| Formation of DNA adducts by ellipticine and its micellar form in Rats – a Comparative Study   | 145 |
| The site directed mutagenesis of key aminoacids in the heme distal side of an oxygen sensor, YddV, probably converts its character from a O <sub>2</sub> sensing protein to a heme oxygenase enzyme | 148 |
| Inhibitors of cyclin-dependent kinases as new generation of anticancer drugs  | 151 |
| Effect of deep freezing to quality of HEK293 Transfected with CaV 3.1 membrane channel  | 153 |
| Fluorescence study of mixed micelles formation  | 156 |
| Interaction of heavy metals with graphene and iron based particles  | 160 |
| FIA-ED: optimization of method for electrochemical study of doxorubicin   | 163 |
| Flow injection analysis with electrochemical detection for rapid identification of platinum-based cytostatics and platinum chlorides in water   | 166 |
| Electrophoretic behavior of doxorubicin   | 169 |
| Utilization of Electrochemistry for detection of bacteria on a 3D printed flow chip   | 173 |
| Liposomal Transporter with GFP mark for Targeted Binding using a Nucleic Acid Anchor System   | 176 |
| Interaction study of arsenic(III) ions with metallothionein gene (MT2a) fragment assessed by spectrometry and electrochemistry  | 179 |
| Automatic Electrochemical determination of heavy metals and application to a remote-controlled robotic platform orpheus-hope  | 182 |
| Microrna electrochemical detection in connection with specific magnetic separation  | 185 |
| Interaction of metallothionein with CdTe quantum dots studied by electrochemistry   | 189 |
| Physical chemistry of inorganic-organic nanostructures  | 192 |
| Elimination square wave voltammetry   | 194 |
| Device for dehydration monitoring using potassium concentration in urine measurement  | 197 |
| Semiconductive SnO <sub>2</sub> /MWCNTs gas sensor  | 200 |
| High resolution technics for fabrication of special nanoelectrodes  | 203 |